



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MISCELLANEOUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

28. Proposed by "IAGO"—(The late DR. JAMES MATTESON, DeKalb Center, Illinois.)

If 9 gentlemen, or 15 ladies, will eat 17 apples in 5 hours, and 15 gentlemen and 15 ladies can eat 47 apples of a similar size in 12 hours, the apples growing uniformly; how many boys will eat up 360 apples in 60 hours, admitting that 120 boys can eat the same number as 18 gentlemen and 26 ladies? *F. P. Matz.*

Solution by Professor P. S. BERG, Larimore, North Dakota.

1. Call the original size of an apple an "apple unit."
2. Call the growth of 1 apple in 1 hour one "unit of growth."
3. $32\frac{1}{4}$ boys will in the same time eat as many as 9 gentlemen or 15 ladies.
4. $\therefore 32\frac{1}{4}$ boys in 5 hours or $160\frac{1}{4}$ boys in 1 hour = 17 "apple units" + 85 "units of growth."
5. $\therefore 85\frac{1}{4}$ boys in 12 hours or $1028\frac{1}{4}$ boys in 1 hour = 47 "apple units" + 564 "units of growth."
6. $\therefore (4) \times 2\frac{1}{4} = 44\frac{1}{4}$ boys in 1 hour = 47 "apple units" + 235 "units of growth."
7. $\therefore (4) \times 6\frac{1}{4} = 106\frac{1}{4}$ boys in 1 hour = $112\frac{1}{4}$ "apple units" + 564 "units of growth."
8. $\therefore (5) - (6) = 584\frac{1}{4}$ boys in 1 hour = 329 "units of growth."
9. $\therefore (7) - (5) = 37\frac{1}{4}$ boys in 1 hour = $65\frac{1}{4}$ "apple units."
10. $\therefore (9) \times 15\frac{1}{4} = 584\frac{1}{4}$ boys in 1 hour = $1016\frac{1}{4}$ "apple units."
11. $\therefore 329$ "units growth" = $1016\frac{1}{4}$ "apple units,"
1 "unit growth" = $3\frac{1}{4}$ "apple units."
12. 360 apples in 60 hours = 360 "apple units" + 21600 "units growth."
13. 360 "apple units" = $116\frac{5}{8}$ "units growth."
14. $\therefore 360$ apples in 60 hours = $116\frac{5}{8} + 21600 = 21716\frac{5}{8}$ "units growth."
15. From (8) 1 boy in 1 hour = $\frac{119 \times 329}{69525}$ "units growth."
16. $\therefore 1$ boy in 60 hours = $1\frac{5}{8} \frac{9}{8}$ "units growth."
17. $\therefore 21716\frac{5}{8} \div 1\frac{5}{8} \frac{9}{8} = 643$ boys.

29. Proposed by ALEXANDER MACFARLANE, M. A., D. Sc., LL. D., Professor of Electrical Engineering in Lehigh University, South Bethlehem, Pennsylvania.

A rectangular room has the four walls, the floor, and the ceiling covered with mirrors; a candle is placed inside the room: find a formula which will express all the images.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

Regard the candle as a luminous point. Then, since there are three sets of parallel mirrors, we have, from elementary optics, three sets of an infinite

number of images situated on three straight lines at right angles to one another, and intersecting at the bright point.

These mirrors are also inclined to one another at right angles. Let $A^\circ = \frac{\pi}{2}$ = the angle of inclination of the mirrors, a° , b° the angles made by the candle with two of the mirrors.

$$\text{Then } \frac{360^\circ - (a^\circ + b^\circ)}{A^\circ} = \frac{360^\circ - 90^\circ}{90^\circ} = 3 = \text{the number of images due to two}$$

of the mirrors inclined at 90° . There are twelve such sets of inclined mirrors, but of the 36 images formed, 18 are repeated. $\therefore \frac{1}{2}$ of 12 of 3 = 18 images due to the inclined mirrors.

$$\therefore \frac{12 \{ 2\pi - (a^\circ + b^\circ) \}}{\pi}, \text{ is the formula for the images due to the inclined}$$

mirrors, where $a^\circ + b^\circ = \frac{\pi}{2}$.

30. Proposed by R. J. ADCOCK, Larchland, Warren County, Illinois.

When the sum of the distances of a point of a plane surface, from all the other points, is a minimum, that point is the center of gravity of the plane surface.

I. Proof by G. B. M. ZERR, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

Let (x_1, y_1) be the point, (x, y) any other point, S the sum of the distances of (x, y) from (x_1, y_1) .

$$\text{Then } S = \int_{x_1}^{x_2} \int_{y_1}^{y_2} \sqrt{(x-x_1)^2 + (y-y_1)^2} dx dy.$$

$$\text{Let } \int_{x_1}^{x_2} \int_{y_1}^{y_2} \text{ be represented by } \int, \text{ and } dx dy \text{ by } dA.$$

$$\therefore S = \int \sqrt{(x-x_1)^2 + (y-y_1)^2} dA = \int D dA.$$

$$\text{For a minimum, } \frac{dS}{dx_1} = \frac{(x-x_1)dA}{D} = 0, \frac{dS}{dy_1} = \frac{(y-y_1)dA}{D} = 0.$$

$$\therefore (x-x_1)dA = 0, (y-y_1)dA = 0. \therefore x_1 = \frac{\int x dA}{\int dA}, y_1 = \frac{\int y dA}{\int dA}.$$

$$\therefore x_1 = \frac{\int \int x dx dy}{\int \int dx dy}, y_1 = \frac{\int \int y dx dy}{\int \int dx dy}.$$